

<2008 英文論文>

Usami M, Kishimoto K, Ohata A, Miyoshi M, Aoyama M, Fueda Y, Kotani J: Butyrate and trichostatin A attenuate nuclear factor κ B activation and tumor necrosis factor- α secretion and increase prostaglandin E2 secretion in peripheral blood mononuclear cells. Nutr Res 28, 321-328, 2008

Taru C, Tuto A, Nakawatase Y, Usami M, Miyawaki I: Gender differences of dietary self-management behavior affecting control indices in type II diabetes. Kobe J Med Sci 54, E82-E96, 2008

Miyoshi M, Usami M, Ohata A: Short-chain fatty acids and trichostatin A alter tight junction permeability in human umbilical vein endothelial cells. Nutrition 24, 1189-1198, 2008

Hirata J, Kotani J, Aoyama M, Kashiwamura S, Ueda H, Kuroda Y, Usami M, Okamura H, Marukawa S: A role for IL-18 in human neutrophil apoptosis. Shock 30, 628-633, 2008

Fujii H1, Hamada Y, Fukagawa M: Bone formation in spontaneously diabetic Torii-newly established model of non-obese type 2 diabetes rats. Bone 42, 372-379, 2008

Hamada Y, Kono TN, Moriguchi Y, Higuchi M, Fukagawa M: Alteration of mRNA expression of molecules related to iron metabolism in adenine-induced renal failure rats: a possible mechanism of iron deficiency in chronic kidney disease patients on treatment. Nephrol Dial Transplant 6, 1886-1891, 2008

Tsuchihashi D, Abe T, Komaba H, Fujii H, Hamada Y, Nii-Kono T, Tanaka M, Fukagawa M: Serum pro-hepcidin as an indicator of iron status in dialysis patients. Ther Apher Dial 12, 226-231, 2008